## Geologic Mapping of Near-Surface Sediments in the Northern Mississippi Embayment

Prepared by Kentucky Research Consortium for Energy and Environment 233 Mining and Minerals Building University of Kentucky, Lexington, KY 40506-0107

Prepared for United States Department of Energy Portsmouth/Paducah Project Office Acknowledgment: This material is based upon work supported by the Department of Energy under Award Number DE-FG05-03OR23032.





April 2005

UK/KRCEE Doc #: P4.2 2005

## Geologic Mapping of Near-Surface Sediments in the Northern Mississippi Embayment


Presented by Josh Sexton Poster Session, Geological Society of America Knoxville, TN

April 2005



## Abstract:


The Jackson Purchase region of western Kentucky consists of Coastal Plain sediments near the northern margin of the Mississippi Embayment. Within this region is the Paducah Gaseous Diffusion Plant (PGDP), a uranium enrichment facility operated by the US Department of Energy. At PGDP, a Superfund site, soil and groundwater studies have provided subsurface lithologic data from hundreds of monitoring wells and borings. Despite preliminary efforts by various contractors, these data have not been utilized to develop detailed stratigraphic correlations of sedimentary units across the study area. In addition, sedimentary exposures along streams in the vicinity of PGDP have not been systematically described beyond the relatively simple geologic quadrangle maps published by the US Geological Survey in 1966–67. Our study integrates lithologic logs, other previous site-investigation data, and outcrop mapping to provide a near-surface stratigraphic framework model of the PGDP area. Preliminary findings suggest that previous ideas about gross heterogeneity of the Holocene and Pleistocene sediments composing the Regional Gravel Aquifer of the area may not be correct. Moreover, the lateral continuity of the Mounds Gravels and the Metropolis Formation may be greater than previously described. These findings may result in a more realistic model of groundwater flow beneath the site. In addition, the findings may be useful for assessing local amplification of earthquake ground motion in near-surface sediments.




Geologic mapping of near-surface sediments in the northern Mississippi Embayment, McCracken County, KY SEXTON, Joshua L., and FRYAR, Alan E., Department of Earth & Environmental Sciences, University of Kentucky, 101 Slone Building, Lexington, KY 40506-0053, jlsexton@uky.edu, and GREB, Stephen F., Kentucky Geological Survey, University of Kentucky, 228 Mining and Mineral Resources Building, Lexington, KY 40506-0107

# Methods

- Formed a digital database compiling lithologic logs from borings and wells installed by PGDP, Tennessee Valley Authority (TVA), UK, and USGS
- Investigated surficial geology where exposures could be found, dominantly along creeks and borrow pits (Site map)
- Documented exposures by measuring sections of outcrop, noting geology, and compiling a photographic database (Image 1)
- Collected sediment samples for geologic age dating using Optically Stimulated Luminescence (OSL) (I mage 2)
- Collected clast orientation data alone exposures where a fabric was observed
- Surveyed contact elevations of stratigraphic units cropping out along Little Bayou Creek and Bayou Creek (I mage 3)
- Examined closely spaced, shallow sediment cores collected during a paleo-seismic investigation (Site map)
- Conducted a field trip with regional experts to discuss site geology and depositional history
- Created stratigraphic cross-sections, isolith maps, and isopach maps of the study area



- Continental Deposits and Quaternary silt and sand at the site



Measured section location along Little Bayou Creek



Image 3: Surveying elevation of Mounds Gravel contact along Little Bayou Creek



Image 2: OSL sampling location along Little Bayou Creek



Image 4: Paleo-liquefaction feature noted along Bayou Creek

- The Mounds Gravel and Metropolis Formation names can be applied respectively to the informal unit names

- Mounds Gravel exposed along Little Bayou Creek and Bayou Creek in the northern portion of the study area were previously unmapped and may be pathways for groundwater discharge to the creeks

- Mounds Gravel appear to form a continuous sheet deposit north of the Ancestral Tennessee River terrace slope interpreted by Jacobs (1997), with areas of channel scour that could have created preferential groundwater flow

- The Metropolis Formation appears to be laterally traceable on the order of 10 to 1000 feet, but does not exhibit traceability beyond this scale because of the small order bedforms associated with a sluggish fluvial depositional system

- Seismic deformation of Quaternary sediments in the study area appears possible from observations noted in previous site investigations, surficial geologic mapping (Image 4), borehole stratigraphic unconformities, and evidence presented in unpublished maps of the Joppa 7.5-minute quadrangle produced by the Illinois State Geological Survey

Jacobs, 1997, Ground-Water Conceptual Model for the Paducah Gaseous Diffusion Plant, Paducah, Kentucky, Prepared for the U.S. Department of Energy under contract DE-AC05-930R22028, Kevil, KY, Jacobs EM Team.

Nelson, W.J., Denny, F. B., Devera, J.A., Follmer, L. R., Masters, J. M., 1999, Quaternary grabens in southernmost Illinois: deformation near an active interplate seismic zone, Tectonophysics, 305, p. 381 - 397.